

**SAGINAW VALLEY  
MEDICAL CONTROL AUTHORITY  
PRACTICE PARAMETER**

**PEDIATRIC  
NAUSEA/VOMITING**

**FIRST RESPONDER**

**I. Management**

- A. Establish and maintain the airway, provide oxygenation and support ventilation.
- B. Position patient on either side if vomiting; otherwise supine.
- C. Obtain vital signs (blood pressure, pulse and respiratory rate).
- D. Do not allow patient to take anything by mouth.
- E. Arrange for ALS intercept.

**BASIC LIFE SUPPORT**

- F. Initiate transport towards ALS intercept, if transporting unit.

**NOTIFY RECEIVING FACILITY**

**ADVANCED LIFE SUPPORT**

- G. Monitor EKG
- H. Consider **IV/IO NORMAL SALINE @ TKO** depending on the clinical circumstances and the results of the EKG monitor.
- I. Consider **ZOFRAN**
  - 1. **IV/IO.1 MG/KG TO A MAXIMUM OF 4MG SLOW OVER 2 MINUTES**
  - 2. **IM= 0.1 MG/KG TO A MAXIMUM OF 4MG**

**CONTACT MEDICAL CONTROL**

MDCIS/EMSCC Approved on: \_\_\_\_\_  
 MCB/MCA Approved on: 17OCT08  
 Supersedes Version Dated: 11DEC07  
 Effective Date: \_\_\_\_\_  
 Revised on: 13OCT08

\_\_\_\_\_  
Medical Director

\_\_\_\_\_  
EMS Manager